

PATENT COOPERATION T EATY

PCT/EP97/0

PCT

NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

United States Patent and Trademark
Office
(Box PCT)
Crystal Plaza 2
Washington, DC 20231
ETATS-UNIS D'AMERIQUE

in its capacity as elected Office

Date of mailing (day/month/year) 05 June 1998 (05.06.98)	Applicant's or agent's file reference 972076wo Me/S
International application No. PCT/EP97/05619	Priority date (day/month/year) 12 October 1996 (12.10.96)
International filing date (day/month/year) 11 October 1997 (11.10.97)	
Applicant KASK, Peet	

1. The designated Office is hereby notified of its election made:

☒ in the demand filed with the International Preliminary Examining Authority on:
09 May 1998 (09.05.98)

☐ in a notice effecting later election filed with the International Bureau on:

2. The election ☒ was
☐ was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

RECEIVED
GROUP 1300
SEP -9 AM 7:22

FEB 05 1999

GROUP 1800

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland Facsimile No.: (41-22) 740.14.35 Form PCT/IB/331 (July 1992)	Authorized officer M. Fourne-Godbersen Telephone No.: (41-22) 338.83.38
---	--

2068648

941220325

17E1

1744

EO/US
PCT/EP97/05639

PATENT COOPERATION TREATY

PCT

NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

United States Patent and Trademark
Office
(Box PCT)
Crystal Plaza 2
Washington, DC 20231
ETATS-UNIS D'AMERIQUE

in its capacity as elected Office

Date of mailing:

23 April 1998 (23.04.98)

International application No.:

PCT/EP97/05639

Applicant's or agent's file reference:

H-21090/A

International filing date:

14 October 1997 (14.10.97)

Priority date:

16 October 1996 (16.10.96)

Applicant:

KARRER, Friedrich et al

1. The designated Office is hereby notified of its election made:



in the demand filed with the International preliminary Examining Authority on:

02 April 1998 (02.04.98)



in a notice effecting later election filed with the International Bureau on:

2. The election ☒ was



was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

98 SEP -9 AM 8:02
RECEIVED
GROUP 1200

RECEIVED

JAN 15 1999

RECEIVED
SERVICE CENTER

The International Bureau of WIPO
34, chemin des Colombettes
1211 Geneva 20, Switzerland

Facsimile No.: (41-22) 740.14.35

Authorized officer:

J. Zahra

Telephone No.: (41-22) 338.83.38

PATENT COOPERATION TREATY

PCT

NOTIFICATION OF THE RECORDING
OF A CHANGE(PCT Rule 92bis.1 and
Administrative Instructions, Section 422)

From the INTERNATIONAL BUREAU

To:

MEYERS, Hans-Wilhelm
Postfach 10 22 41
D-50462 Köln
ALLEMAGNEDate of mailing (day/month/year)
11 December 1998 (11.12.98)Applicant's or agent's file reference
972076wo Me/S

IMPORTANT NOTIFICATION

International application No.
PCT/EP97/05619International filing date (day/month/year)
11 October 1997 (11.10.97)

1. The following indications appeared on record concerning:

☒ the applicant ☐ the inventor ☐ the agent ☐ the common representative

Name and Address

EVOTEC BIOSYSTEMS GMBH
Grandweg 64
D-22529 Hamburg
Germany

State of Nationality

DE

State of Residence

DE

Telephone No.

Facsimile No.

Teleprinter No.

2. The International Bureau hereby notifies the applicant that the following change has been recorded concerning:

☐ the person ☒ the name ☒ the address ☐ the nationality ☐ the residence

Name and Address

EVOTEC BIOSYSTEMS AG
Schnackenburgallee 114
D-22525 Hamburg
Germany

State of Nationality

DE

State of Residence

DE

Telephone No.

Facsimile No.

Teleprinter No.

3. Further observations, if necessary:

4. A copy of this notification has been sent to:

☒ the receiving Office ☐ the designated Offices concerned
☐ the International Searching Authority ☒ the elected Offices concerned
☒ the International Preliminary Examining Authority ☐ other:The International Bureau of WIPO
34, chemin des Colombettes
1211 Geneva 20, Switzerland

Authorized officer

G. Bähr

Facsimile No.: (41-22) 740.14.35

Telephone No.: (41-22) 338.83.38

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : G01N 21/64	A1	(11) International Publication Number: WO 98/57150
		(43) International Publication Date: 17 December 1998 (17.12.98)

(21) International Application Number: PCT/EP98/03509
(22) International Filing Date: 10 June 1998 (10.06.98)

(30) Priority Data:
97109353.9 10 June 1997 (10.06.97) EP
(34) Countries for which the regional or
international application was filed: DE et al.

(71) Applicant (for all designated States except US): EVOTEC
BIOSYSTEMS GMBH [DE/DE]; Schnackenburgallee 114,
D-22525 Hamburg (DE).

(72) Inventor; and
(75) Inventor/Applicant (for US only): KASK, Peet [EE/EE]; Uus
5-3, EE-3051 Harku (EE).

(74) Agents: MEYERS, Hans-Wilhelm et al.; P.O. Box 10 22 41,
D-50462 Köln (DE).


(81) Designated States: JP, US, European patent (AT, BE, CH, CY,
DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT,
SE).

Published

With international search report.

*Before the expiration of the time limit for amending the
claims and to be republished in the event of the receipt of
amendments.*

(54) Title: A METHOD FOR CHARACTERIZING SAMPLES ON THE BASIS OF INTERMEDIATE STATISTICAL DATA

**(57) Abstract**

A method for characterizing samples which contain units, comprising the steps of: a) monitoring intensity fluctuations of radiation emitted, scattered and/or reflected by the units in at least one measurement volume with at least one detection means which is capable of detecting radiation emitted, scattered and/or reflected by said units, b) determining from said intensity fluctuations intermediate statistical data comprising an at least two-dimensional joint statistical function, c) determining information related to a joint distribution of units out of said intermediate statistical data.

CLAIMS

1. A method for characterizing samples which contain units, comprising the steps of:
 - a) monitoring intensity fluctuations of radiation emitted, scattered and/or reflected by the units in at least one measurement volume with at least one detection means which is capable to detect radiation emitted, scattered and/or reflected by said units,
 - b) determining from said intensity fluctuations intermediate statistical data comprising an at least two-dimensional joint statistical function,
 - c) determining information related to a joint distribution of units out of said intermediate statistical data.
2. The method according to claim 1, wherein in step c) the presence or absence or concentration of at least one unit or species of units with a specific combination of at least two physical properties is determined.
3. The method according to claim 1 and/or 2, wherein in step c) a joint distribution of units as a function of at least two specific physical properties is determined out of said intermediate statistical data.
4. The method according to at least one of claims 1 to 3, wherein in step c) a function of said joint distribution is determined.
5. The method according to at least one of claims 1 to 4, wherein the intermediate statistical data in step b) are derived from the intensity fluctuations directly or from fluctuations of local statistical functions of intensity.
6. The method according to at least one of claims 1 to 5, wherein said intensity

fluctuations are monitored in terms of determining numbers of photon counts in, preferably consecutive, time intervals of given length.

7. The method according to at least one of claims 1 to 6, wherein said intensity fluctuations are monitored in terms of determining time of arrival of photons and/or length of time intervals between a given number of consecutive photon counts.
8. The method according to at least one of claims 1 to 7, wherein said intermediate statistical data additionally comprise one-dimensional statistical functions.
9. The method according to at least one of claims 1 to 8, wherein at least one of the arguments of said joint statistical function is time.
10. The method according to at least one of claims 1 to 9, wherein at least one of the arguments of said joint statistical function is the number of photon counts in time intervals of given length or length of time intervals between a given number of consecutive photon counts.
11. The method according to at least one of claims 1 to 10, wherein at least one of the arguments of said joint statistical function belongs to the class of arguments of mathematical transforms of either probability or correlation functions.
12. The method according to at least one of the claims 1 to 11, wherein said intermediate statistical data are selected from a group consisting of unconditional and conditional distributions of the number of photon counts, distributions of the length of time intervals between photon counts, autocorrelation functions, cross-correlation functions, and combinations thereof.
13. The method according to at least one of the claims 1 to 12, wherein said units are particles, molecules, aggregates, vesicles, cells, viruses, bacteria, beads, centers, or mixtures thereof in solids, liquids or gases.

14. The method according to at least one of the claims 1 to 13, wherein said units can be grouped into species which can be distinguished by at least one of their specific physical properties.
15. The method according to at least one of the claims 1 to 14, wherein at least one species is luminescent, preferably fluorescent, and/or is luminescently labelled and at least one detection means monitors the fluctuations of luminescence intensity.
16. The method according to at least one of the claims 1 to 15, wherein at least one of the specific physical properties characterizing said units is the diffusion coefficient, or correlation time of radiation intensity fluctuations, or any other property directly related to said diffusion coefficient.
17. The method according to at least one of the claims 1 to 16, wherein at least one of the specific physical properties characterizing said units is the specific brightness.
18. The method according to at least one of the claims 1 to 17, wherein at least one of the specific physical properties characterizing fluorescent units is the polarization ratio of their fluorescence, or fluorescence anisotropy, or any other property expressing the extent of polarization of fluorescence.
19. The method according to at least one of the claims 1 to 18, wherein at least one of the specific physical properties characterizing fluorescent units is the ratio of fluorescence intensities corresponding to different excitation wavelengths and/or different spectral sensitivities of fluorescence detection, or any other property expressing the dependence of fluorescence intensity on the wavelength of excitation and/or detection.
20. The method according to at least one of the claims 1 to 19, wherein at least one of the specific physical properties characterizing fluorescent units is lifetime of fluorescence.

21. The method according to at least one of the claims 1 to 20, wherein the specific physical properties, in particular luminescence properties like fluorescence lifetime or fluorescence anisotropy, of the units are varied by conjugating them with a specific luminophore via different linker molecules.
22. The method according to at least one of the claims 1 to 21, wherein the luminescence properties of the units are varied by conjugating them with a first molecule, in particular biotin, which binds a luminescently labelled second molecule, in particular luminescently labelled avidin or streptavidin.
23. The method according to at least one of the claims 1 to 22, wherein the luminescence properties of a unit are changed by energy transfer, in which energy absorbed by said unit is transferred upon close contact to a luminophore of an acceptor unit and subsequently emitted.
24. The method according to at least one of the claims 1 to 23 for use in high throughput screening, diagnostics, monitoring of polymerisation, aggregation and degradation processes, particle sorting, or nucleic acid sequencing.
25. The method according to at least one of the claims 1 to 24, wherein said intermediate statistical data are fitted using a priori information on said sample.
26. The method according to at least one of the claims 1 to 25, wherein said statistical data are processed applying multidimensional inverse transformation with linear regularization and/or constraints.
27. The method according to at least one of the claims 1 to 26, wherein the measurement volume is only a part of the total volume of said sample and has a volume $\leq 10^{-12}$ l, preferably $\leq 10^{-14}$ l.

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 98/03509

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 G01N21/64

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0 359 681 A (UNIVERSITY OF ARKANSAS) 21 March 1990 see page 2, line 3 - line 9 see page 5, line 38 - line 40 see page 8, line 2 - line 6 see page 9, line 14 - line 23 see page 10, line 48 - line 54 ---	1,7,8, 12,13, 15,16,24
Y	US 3 824 402 A (MULLANEY) 16 July 1974 see abstract see column 3, line 58 - column 4, line 21 --- -/--	1,7,8, 12,13, 15,16,24



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

20 October 1998

Date of mailing of the international search report

28/10/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer:

Thomas, R.M.

PCT

INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference 972076wo Me/S	FOR FURTHER ACTION see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.	
International application No. PCT/EP 97/05619	International filing date (day/month/year) 11/10/1997	(Earliest) Priority Date (day/month/year) 12/10/1996
Applicant EVOTEC BIOSYSTEMS GMBH et al.		

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 3 sheets.

☒ It is also accompanied by a copy of each prior art document cited in this report.

1. ☐ **Certain claims were found unsearchable** (see Box I).
2. ☐ **Unity of invention is lacking** (see Box II).
3. ☐ The international application contains disclosure of a **nucleotide and/or amino acid sequence listing** and the international search was carried out on the basis of the sequence listing
 - ☐ filed with the international application.
 - ☐ furnished by the applicant separately from the international application.
 - ☐ but not accompanied by a statement to the effect that it did not include matter going beyond the disclosure in the international application as filed.
 - ☐ Transcribed by this Authority
4. With regard to the **title**, ☒ the text is approved as submitted by the applicant
☐ the text has been established by this Authority to read as follows:
5. With regard to the **abstract**, ☒ the text is approved as submitted by the applicant
☐ the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this International Search Report, submit comments to this Authority.
6. The figure of the **drawings** to be published with the abstract is:
 Figure No. 1 ☒ as suggested by the applicant. ☐ None of the figures.
☐ because the applicant failed to suggest a figure.
☐ because this figure better characterizes the invention.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 97/05619

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 G01N15/02

According to International Patent Classification(IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	H. QIAN ET AL.: "Distribution of molecular aggregation by analysis of fluctuation moments" PROC. NATL. ACAD. SCI. , vol. 87, July 1990, USA, pages 5479-5483, XP000647943 cited in the application see page 5479, column 1, paragraph 1 see page 5479, column 2, paragraph 3 see figure 3 see page 5483, column 1, paragraph 2 ----- -/--	1

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

° Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

16 February 1998

Date of mailing of the international search report

24/02/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Zinngrebe, U

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 97/05619

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	SM. NIE ET AL.: "real-time detection of single-molecules in solution by confocal fluorescence microscopy" ANALYTICAL CHEMISTRY, vol. 67, no. 17, 1 September 1995, COLUMBUS US, pages 2849-2857, XP000647270 see figure 3 ----	1
A	EP 0 601 714 A (HAMAMATSU PHOTONICS KK) 15 June 1994 see page 3, line 19-27 see page 5, line 5-26 see page 9, line 55 - page 10, line 51 see page 12, line 3-21 ----	1
A	US 4 788 443 A (FURUYA YOSHIYUKI) 29 November 1988 see column 1, line 5-23 see column 3, line 30 - column 4 see figures 1,4 ----	1
A	US 4 768 879 A (MCLACHLAN RICHARD D ET AL) 6 September 1988 see abstract see column 4, line 59 - column 5, line 15; figure 2 -----	1

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 97/05619

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0601714 A	15-06-94	JP 2575270 B	22-01-97
		JP 6148076 A	27-05-94
		US 5528046 A	18-06-96

US 4788443 A	29-11-88	JP 1878140 C	07-10-94
		JP 6003413 B	12-01-94
		JP 62287130 A	14-12-87

US 4768879 A	06-09-88	NONE	

28 Rec'd PCT/PTO 10 MAR 1998
PCT

ANTRAG

Der Unterzeichnete beantragt, daß die vorliegende internationale Anmeldung nach dem Vertrag über die internationale Zusammenarbeit auf dem Gebiet des Patentwesens behandelt wird.

Vom Anmeldeamt auszufüllen

Internationales Aktenzeichen

Internationales Anmeldedatum

Name des Anmeldeamts und "PCT International Application"

Aktenzeichen des Anmelders oder Anwalts (falls gewünscht)
(max. 12 Zeichen) 972076wo Me/Sch-gn

Feld Nr. I BEZEICHNUNG DER ERFINDUNG Method of analysis of samples by determination of a function of specific brightness

Feld Nr. II ANMELDER

Name und Anschrift: (Familienname, Vorname; bei juristischen Personen vollständige amtliche Bezeichnung. Bei der Anschrift sind die Postleitzahl und der Name des Staats anzugeben.)

EVOTEC BIOSYSTEMS GMBH
Grandweg 64

22529 Hamburg
DE

☐ Diese Person ist gleichzeitig Erfinder

Telefonnr.:

Telefaxnr.:

Fernschreibnr.:

Staatsangehörigkeit (Staat): DE

Sitz oder Wohnsitz (Staat): DE

Diese Person ist Anmelder für folgende Staaten:

☐ alle Bestimmungsstaaten

☒ alle Bestimmungsstaaten mit Ausnahme der Vereinigten Staaten von Amerika

☐ nur die Vereinigten Staaten von Amerika

☐ die im Zusatzfeld angegebenen Staaten

Feld Nr. III WEITERE ANMELDER UND/ODER (WEITERE) ERFINDER

Name und Anschrift: (Familienname, Vorname; bei juristischen Personen vollständige amtliche Bezeichnung. Bei der Anschrift sind die Postleitzahl und der Name des Staats anzugeben.)

KASK, Pete
Kirsi 8-104, Tallinn

EE0006 Estonia

Diese Person ist:

☐ nur Anmelder

☐ Anmelder und Erfinder

☐ nur Erfinder (Wird dieses Kästchen angekreuzt, so sind die nachstehenden Angaben nicht nötig.)

Staatsangehörigkeit (Staat): EE

Sitz oder Wohnsitz (Staat): EE

Diese Person ist Anmelder für folgende Staaten:

☐ alle Bestimmungsstaaten

☐ alle Bestimmungsstaaten mit Ausnahme der Vereinigten Staaten von Amerika

☒ nur die Vereinigten Staaten von Amerika

☐ die im Zusatzfeld angegebenen Staaten

☐ Weitere Anmelder und/oder (weitere) Erfinder sind auf einem Fortsetzungsblatt angegeben.

Feld Nr. IV ANWALT ODER GEMEINSAMER VERTRETER; ZUSTELLANSCHRIFT

Die folgende Person wird hiermit bestellt/ist bestellt worden, um für den (die) Anmelder vor den zuständigen internationalen Behörden in folgender Eigenschaft zu handeln als: ☒ Anwalt ☐ gemeinsamer Vertreter

Name und Anschrift: (Familienname, Vorname; bei juristischen Personen vollständige amtliche Bezeichnung. Bei der Anschrift sind die Postleitzahl und der Name des Staats anzugeben.)

MEYERS, Hans-Wilhelm
Postfach 10 22 41

50462 Köln
DE

Telefonnr.:
0221 916520

Telefaxnr.:
0221 134297

Fernschreibnr.:

☐ Dieses Kästchen ist anzukreuzen, wenn kein Anwalt oder gemeinsamer Vertreter bestellt ist und statt dessen im obigen Feld eine spezielle Zustellanschrift angegeben ist.

Feld Nr. V BESTIMMUNG VON STAATEN

Die folgenden Bestimmungen nach Regel 4.9 Absatz a werden hiermit vorgenommen (bitte die entsprechenden Kästchen ankreuzen; wenigstens ein Kästchen muß angekreuzt werden):

Regionales Patent

- ☒ **AP ARIPO-Patent:** GH Ghana, KE Kenia, LS Lesotho, MW Malawi, SD Sudan, SZ Swasiland, UG Uganda, ZW Simbabwe und jeder weitere Staat, der Vertragsstaat des Harare-Protokolls und des PCT ist
- ☒ **EA Eurasisches Patent:** AM Armenien, AZ Aserbaidshan, BY Belarus, KG Kirgisistan, KZ Kasachstan, MD Republik Moldau, RU Russische Föderation, TJ Tadschikistan, TM Turkmenistan und jeder weitere Staat, der Vertragsstaat des Eurasischen Patentübereinkommens und des PCT ist
- ☒ **EP Europäisches Patent:** AT Österreich, BE Belgien, CH und LI Schweiz und Liechtenstein, DE Deutschland, DK Dänemark, ES Spanien, FI Finnland, FR Frankreich, GB Vereinigtes Königreich, GR Griechenland, IE Irland, IT Italien, LU Luxemburg, MC Monaco, NL Niederlande, PT Portugal, SE Schweden und jeder weitere Staat, der Vertragsstaat des Europäischen Patentübereinkommens und des PCT ist
- ☒ **OA OAPI-Patent:** BF Burkina Faso, BJ Benin, CF Zentralafrikanische Republik, CG Kongo, CI Côte d'Ivoire, CM Kamerun, GA Gabun, GN Guinea, ML Mali, MR Mauretanien, NE Niger, SN Senegal, TD Tschad, TG Togo und jeder weitere Staat, der Vertragsstaat der OAPI und des PCT ist (falls eine andere Schutzrechtsart oder ein sonstiges Verfahren gewünscht wird, bitte auf der gepunkteten Linie angeben)

Nationales Patent (falls eine andere Schutzrechtsart oder ein sonstiges Verfahren gewünscht wird, bitte auf der gepunkteten Linie angeben):

- | | |
|--|---|
| <input checked="" type="checkbox"/> AL Albanien | <input checked="" type="checkbox"/> LV Lettland |
| <input type="checkbox"/> AM Armenien | <input type="checkbox"/> MD Republik Moldau |
| <input type="checkbox"/> AT Österreich | <input checked="" type="checkbox"/> MG Madagaskar |
| <input checked="" type="checkbox"/> AU Australien | <input checked="" type="checkbox"/> MK Die ehemalige jugoslawische Republik
Mazedonien |
| <input type="checkbox"/> AZ Aserbaidshan | <input checked="" type="checkbox"/> MN Mongolei |
| <input checked="" type="checkbox"/> BA Bosnien-Herzegowina | <input type="checkbox"/> MW Malawi |
| <input checked="" type="checkbox"/> BB Barbados | <input checked="" type="checkbox"/> MX Mexiko |
| <input checked="" type="checkbox"/> BG Bulgarien | <input type="checkbox"/> NO Norwegen |
| <input checked="" type="checkbox"/> BR Brasilien | <input checked="" type="checkbox"/> NZ Neuseeland |
| <input type="checkbox"/> BY Belarus | <input checked="" type="checkbox"/> PL Polen |
| <input checked="" type="checkbox"/> CA Kanada | <input type="checkbox"/> PT Portugal |
| <input type="checkbox"/> CH und LI Schweiz und Liechtenstein | <input checked="" type="checkbox"/> RO Rumänien |
| <input checked="" type="checkbox"/> CN China | <input type="checkbox"/> RU Russische Föderation |
| <input checked="" type="checkbox"/> CU Kuba | <input type="checkbox"/> SD Sudan |
| <input checked="" type="checkbox"/> CZ Tschechische Republik | <input type="checkbox"/> SE Schweden |
| <input type="checkbox"/> DE Deutschland | <input checked="" type="checkbox"/> SG Singapur |
| <input type="checkbox"/> DK Dänemark | <input checked="" type="checkbox"/> SI Slowenien |
| <input checked="" type="checkbox"/> EE Estland | <input checked="" type="checkbox"/> SK Slowakei |
| <input type="checkbox"/> ES Spanien | <input checked="" type="checkbox"/> SL Sierra Leone |
| <input type="checkbox"/> FI Finnland | <input type="checkbox"/> TJ Tadschikistan |
| <input type="checkbox"/> GB Vereinigtes Königreich | <input type="checkbox"/> TM Turkmenistan |
| <input checked="" type="checkbox"/> GE Georgien | <input checked="" type="checkbox"/> TR Türkei |
| <input type="checkbox"/> GH Ghana | <input checked="" type="checkbox"/> TT Trinidad und Tobago |
| <input checked="" type="checkbox"/> HU Ungarn | <input checked="" type="checkbox"/> UA Ukraine |
| <input checked="" type="checkbox"/> IL Israel | <input type="checkbox"/> UG Uganda |
| <input checked="" type="checkbox"/> IS Island | <input checked="" type="checkbox"/> US Vereinigte Staaten von Amerika |
| <input checked="" type="checkbox"/> JP Japan | <input type="checkbox"/> UZ Usbekistan |
| <input type="checkbox"/> KE Kenia | <input checked="" type="checkbox"/> VN Vietnam |
| <input type="checkbox"/> KG Kirgisistan | <input checked="" type="checkbox"/> YU Jugoslawien |
| <input checked="" type="checkbox"/> KP Demokratische Volksrepublik Korea | <input type="checkbox"/> ZW Simbabwe |
| <input checked="" type="checkbox"/> KR Republik Korea | |
| <input type="checkbox"/> KZ Kasachstan | |
| <input checked="" type="checkbox"/> LC Saint Lucia | |
| <input checked="" type="checkbox"/> LK Sri Lanka | |
| <input checked="" type="checkbox"/> LR Liberia | |
| <input type="checkbox"/> LS Lesotho | |
| <input checked="" type="checkbox"/> LT Litauen | |
| <input type="checkbox"/> LU Luxemburg | |

Kästchen für die Bestimmung von Staaten (für die Zwecke eines nationalen Patents), die dem PCT nach der Veröffentlichung dieses Formblatts beigetreten sind:

☒ und jeder weitere Staat

Zusätzlich zu den oben genannten Bestimmungen nimmt der Anmelder nach Regel 4.9 Absatz b auch alle anderen nach dem PCT zulässigen Bestimmungen vor mit Ausnahme der Bestimmung von

Der Anmelder erklärt, daß diese zusätzlichen Bestimmungen unter dem Vorbehalt einer Bestätigung stehen und jede zusätzliche Bestimmung, die vor Ablauf von 15 Monaten ab dem Prioritätsdatum nicht bestätigt wurde, nach Ablauf dieser Frist als vom Anmelder zurückgenommen gilt. (Die Bestätigung einer Bestimmung erfolgt durch die Einreichung einer Mitteilung, in der diese Bestimmung angegeben wird, und die Zahlung der Bestimmungs- und der Bestätigungsgebühr. Die Bestätigung muß beim Anmeldeamt innerhalb der Frist von 15 Monaten eingehten.)

Zusatzfeld Wird dieses Zusatzfeld nicht benutzt, so ist dieses Blatt dem Antrag nicht beizufügen.

Dieses Feld ist in folgenden Fällen auszufüllen:

1. Wenn der Platz in einem Feld nicht für alle Angaben ausreicht:
insbesondere:
 - i) Wenn mehr als zwei Anmelder und/oder Erfinder vorhanden sind und kein Fortsetzungsblatt zur Verfügung steht:
In diesem Fall sind mit dem Vermerk "Fortsetzung von Feld Nr. ..." [Nummer des Feldes angeben] die gleichen Angaben zu machen wie in dem Feld vorgesehen, das platzmäßig nicht ausreicht;
 - ii) Wenn in Feld Nr. II oder III die Angabe "die im Zusatzfeld angegebenen Staaten" angekreuzt ist:
In diesem Fall sind mit dem Vermerk "Fortsetzung von Feld Nr. II", "Fortsetzung von Feld Nr. III" oder "Fortsetzung von Feld Nr. II und Nr. III" die Namen der Anmelder und neben jedem Namen der Staat oder die Staaten (und/oder ggf. ARIPO-, eurasisches, europäisches oder OAPI-Patent) anzugeben, für die die bezeichnete Person Anmelder ist.
 - iii) Wenn der in Feld Nr. II oder III genannte Erfinder oder Erfinder/Anmelder nicht für alle Bestimmungsstaaten oder für die Vereinigten Staaten von Amerika als Erfinder benannt ist:
In diesem Fall sind mit dem Vermerk "Fortsetzung von Feld Nr. II" oder "Fortsetzung von Feld Nr. III" oder "Fortsetzung von Feld Nr. II und Nr. III" der Name des Erfinders und neben jedem Namen der Staat oder die Staaten (und/oder ggf. ARIPO-, eurasisches, europäisches oder OAPI-Patent) anzugeben, für die die bezeichnete Person Erfinder ist.
 - iv) Wenn zusätzlich zu dem Anwalt/den Anwälten, die in Feld Nr. IV angegeben sind, weitere Anwälte bestellt sind:
In diesem Fall sind mit dem Vermerk "Fortsetzung von Feld Nr. IV" für jeden weiteren Anwalt die gleichen Angaben zu machen wie in Feld Nr. IV vorgesehen.
 - v) Wenn in Feld Nr. V bei einem Staat (oder bei OAPI) die Angabe "Zusatzpatent" oder "Zusatzzertifikat" oder wenn in Feld Nr. V bei den Vereinigten Staaten von Amerika die Angabe "Fortsetzung" oder "Teilfortsetzung" hinzugefügt wird:
In diesem Fall sind mit dem Vermerk "Fortsetzung von Feld Nr. V" die Namen der betreffenden Staaten (oder OAPI) und nach dem Namen jeder dieser Staaten (oder OAPI) das Aktenzeichen des Hauptschutzrechts oder der Hauptschutzrechtsanmeldung und das Datum der Erteilung des Hauptschutzrechts oder der Einreichung der Hauptschutzrechtsanmeldung anzugeben.
 - vi) Wenn die Priorität von mehr als drei früheren Anmeldungen beansprucht wird:
In diesem Fall sind mit dem Vermerk "Fortsetzung von Feld Nr. VI" für jede weitere frühere Anmeldung die gleichen Angaben zu machen wie in Feld Nr. VI vorgesehen.
2. Wenn der Anmelder für irgendein Bestimmungsamt die Vergünstigung nationaler Vorschriften betreffend unschädliche Offenbarung oder Ausnahmen von der Neuheitsschädlichkeit in Anspruch nimmt:
In diesem Fall ist mit dem Vermerk "Erklärung betreffend unschädliche Offenbarung oder Ausnahmen von der Neuheitsschädlichkeit" nachstehend diese Erklärung abzugeben.

Fortsetzung zu Feld Nr. IV

VON KREISLER, Alek
 SELTING, Günther
 WERNER, Hans-Karsten
 FUESS, F. Johann
 DALLMEYER, Georg
 HILLERINGMANN, Jochen
 JÖNSSON, Hans-Peter
 WEBER, Thomas
 HELBING, Jörg

Postfach 10 22 41

50462 Köln
 DE

Feld Nr. VI PRIORITÄTSANSPRUCHWeitere Prioritätsansprüche sind im Zusatzfeld angegeben. ☐

Die Priorität der folgenden früheren Anmeldung(en) wird hiermit beansprucht:

Staat (Anmelde- oder Bestimmungsstaat der Anmeldung)	Anmeldedatum (Tag/Monat/Jahr)	Aktenzeichen	Anmeldeamt (nur bei regionaler oder internationaler Anmeldung)
(1) EP (DE)	12. Oktober 1996 (12.10.1996)	96 116 373.0	
(2)			
(3)			

Dieses Kästchen ankreuzen, wenn die beglaubigte Kopie der früheren Anmeldung von dem Amt ausgestellt werden soll, das für die Zwecke dieser internationalen Anmeldung Anmeldeamt ist (eine Gebühr kann verlangt werden):

☒ Das Anmeldeamt wird hiermit ersucht, eine beglaubigte Abschrift der oben in Zeile(n) (1) bezeichneten früheren Anmeldung(en) zu erstellen und dem Internationalen Büro zu übermitteln.
Feld Nr. VII INTERNATIONALE RECHERCHENBEHÖRDE

Wahl der Internationalen Recherchenbehörde (ISA) (Sind zwei oder mehr Internationale Recherchenbehörden für die internationale Recherche zuständig, ist der Name der Behörde anzugeben, die die internationale Recherche durchführen soll; Zweibuchstaben-Code genügt):

ISA /

Frühere Recherche: Auszufüllen, wenn eine Recherche (internationale Recherche, Recherche internationaler Art oder sonstige Recherche) bereits bei der internationalen Recherchenbehörde beantragt oder von ihr durchgeführt worden ist und diese Behörde nun ersucht wird, die internationale Recherche soweit wie möglich auf die Ergebnisse einer solchen früheren Recherche zu stützen. Die Recherche oder der Recherchenantrag ist durch Angabe der betreffenden Anmeldung (bzw. deren Übersetzung) oder des Recherchenantrags zu bezeichnen.

Staat (oder regionales Amt):

EPA

Datum (Tag/Monat/Jahr):

15. April 1997

Aktenzeichen:

96 116 373.0

Feld Nr. VIII KONTROLLISTE

Diese internationale Anmeldung umfasst:

1. Antrag : 4 Blätter
 2. Beschreibung : 27 Blätter
 3. Ansprüche : 5 Blätter
 4. Zusammenfassung : 1 Blätter
 5. Zeichnungen : 12 Blätter
 Insgesamt : 49 Blätter

Dieser internationalen Anmeldung liegen die nachstehend angekreuzten Unterlagen bei:

1. ☐ Unterzeichnete gesonderte Vollmacht
 2. ☐ Kopie der allgemeinen Vollmacht
 3. ☐ Begründung für das Fehlen der Unterschrift
 4. ☐ Prioritätsbeleg(e) (durch die Zeilennummer von Feld Nr. VI kennzeichnen):
 5. ☒ Blatt für die Gebührenberechnung
 6. ☐ Gesonderte Angaben zu hinterlegten Mikroorganismen
 7. ☐ Sequenzprotokolle für Nucleotide und/oder Aminosäuren (Diskette)
 8. ☒ Sonstige (einzeln auflisten):
 EP 96 116 373.0

Abbildung Nr. 1 der Zeichnungen (falls vorhanden) soll mit der Zusammenfassung veröffentlicht werden.**Feld Nr. IX UNTERSCHRIFT DES ANMELDERS ODER DES ANWALTS**

Der Name jeder unterzeichnenden Person ist neben der Unterschrift zu wiederholen, und es ist anzugeben, sofern sich dies nicht eindeutig aus dem Antrag ergibt, in welcher Eigenschaft die Person unterzeichnet.

Der Patentanwalt

(Dr. Meyers)

Vom Anmeldeamt auszufüllen

1. Datum des tatsächlichen Eingangs dieser internationalen Anmeldung:

3. Geändertes Eingangsdatum aufgrund nachträglich, jedoch fristgerecht eingegangener Unterlagen oder Zeichnungen zur Vervollständigung dieser internationalen Anmeldung:

4. Datum des fristgerechten Eingangs der angeforderten Richtigstellungen nach Artikel 11(2) PCT:

5. Vom Anmelder benannte Internationale Recherchenbehörde:

ISA /

6. ☐ Übermittlung des Recherchenexemplars bis zur Zahlung der Recherchegebühr aufgeschoben

2. Zeichnungen

☐ eingegangen:☐ nicht eingegangen:

Vom Internationalen Büro auszufüllen

Datum des Eingangs des Aktenexemplars beim Internationalen Büro:

PCT

BLATT FÜR DIE GEBÜHRENBERECHNUNG Anhang zum Antrag

Von Anmeldamt auszufüllen

Internationales Aktenzeichen

Eingangsstempel des Anmeldamts

Aktenzeichen des Anmelders

oder Anwalts 972076wo Me/Sch-gn

Anmelder EVOTEC BIOSYSTEMS GMBH, Granweg 64
22529 Hamburg
DE

BERECHNUNG DER VORGESCHRIEBENEN GEBÜHREN

1. ÜBERMITTLUNGSGEBÜHR 200,-- Ü
2. RECHERCHENGEBÜHR 2.200,-- R

Die internationale Recherche ist durchzuführen von
(Sind zwei oder mehr Internationale Recherchebehörden für die internationale Recherche zuständig,
ist der Name der Behörde anzugeben, die die internationale Recherche durchführen soll.)

3. INTERNATIONALE GEBÜHR

Grundgebühr

Die internationale Anmeldung enthält 49 Blätter.

umfaßt die ersten 30 Blätter 955,-- g₁

19 x 19,-- = 361,-- g₂

Anzahl der Blätter Zusatzblattgebühr
über 30

Addieren Sie die in Feld g₁ und g₂ eingetragenen
Beträge, und tragen Sie die Summe in Feld G ein 1.316,-- G

Bestimmungsgebühren

Die internationale Anmeldung enthält alle Bestimmungen.

11 x 232,-- = 2.552,-- B

Anzahl der zu zahlenden Bestimmungsgebühren (maximal 11)

Addieren Sie die in Feld G und B eingetragenen
Beträge, und tragen Sie die Summe in Feld I ein
(Anmelder aus einigen Staaten haben Anspruch auf eine Ermäßigung der internationalen Gebühr von
75%. Hat der Anmelder (oder haben alle Anmelder) einen solchen Anspruch, so beträgt der in Feld I
einzutragende Gesamtbetrag 25% der Summe der in Feld G und B eingetragenen Beträge.)

3.868,-- I

4. GEBÜHR FÜR PRIORITÄTSBELEG

GESAMTBETRAG DER ZU ZAHLENDEN GEBÜHREN

Addieren Sie die in Feldern Ü, R, I und P eingetragenen Beträge,
und tragen Sie die Summe in das nebenstehende Feld ein

60,-- P

6.328,--

INSGESAMT

☐ Die Bestimmungsgebühren werden jetzt noch nicht gezahlt.

ZAHLUNGSWEISE

- ☒ Abbuchungsauftrag (siehe unten) ☐ Bankwechsel ☐ Kupons
- ☐ Scheck ☐ Barzahlung ☐ Sonstige (einzeln angeben):
- ☐ Postanweisung ☐ Gebührenmarken

ABBUCHUNGSauftrag (diese Zahlungsweise gibt es nicht bei allen Anmeldeämtern)

- Das Anmeldeamt/ ☒ wird beauftragt, den vorstehend angegebenen Gesamtbetrag der Gebühren von meinem laufenden Konto abzubuchen.
- ☐ wird beauftragt, Fehlbeträge oder Überzahlungen des vorstehend angegebenen Gesamtbetrags der Gebühren meinem laufenden Konto zu belasten bzw. gutzuschreiben.
- ☐ wird beauftragt, die Gebühr für die Ausstellung des Prioritätsbelegs und seine Übermittlung an das Internationale Büro der WIPO von meinem laufenden Konto abzubuchen.

2800.0007

10. Oktober 1997

Contonummer

Datum (Tag/Monat/Jahr)

Unterschrift (Dr. Meyers)

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 97/05619

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 G01N15/02

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category,*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	H. QIAN ET AL.: "Distribution of molecular aggregation by analysis of fluctuation moments" PROC. NATL. ACAD. SCI., vol. 87, July 1990, USA, pages 5479-5483, XP000647943 cited in the application see page 5479, column 1, paragraph 1 see page 5479, column 2, paragraph 3 see figure 3 see page 5483, column 1, paragraph 2 --- -/--	1

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

16 February 1998

Date of mailing of the international search report

24/02/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Zinngrebe, U

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 97/05619

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	SM. NIE ET AL.: "real-time detection of single-molecules in solution by confocal fluorescence microscopy" ANALYTICAL CHEMISTRY, vol. 67, no. 17, 1 September 1995, COLUMBUS US, pages 2849-2857, XP000647270 see figure 3 -----	1
A	EP 0 601 714 A (HAMAMATSU PHOTONICS KK) 15 June 1994 see page 3, line 19-27 see page 5, line 5-26 see page 9, line 55 - page 10, line 51 see page 12, line 3-21 -----	1
A	US 4 788 443 A (FURUYA YOSHIYUKI) 29 November 1988 see column 1, line 5-23 see column 3, line 30 - column 4 see figures 1,4 -----	1
A	US 4 768 879 A (MCLACHLAN RICHARD D ET AL) 6 September 1988 see abstract see column 4, line 59 - column 5, line 15; figure 2 -----	1

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 97/05619

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0601714 A	15-06-94	JP 2575270 B	22-01-97
		JP 6148076 A	27-05-94
		US 5528046 A	18-06-96
<hr/>			
US 4788443 A	29-11-88	JP 1878140 C	07-10-94
		JP 6003413 B	12-01-94
		JP 62287130 A	14-12-87
<hr/>			
US 4768879 A	06-09-88	NONE	
<hr/>			

fa 5000

9/029830



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification G01N 15/02		(11) International Publication Number: WO 98/16814
A1		(43) International Publication Date: 23 April 1998 (23.04.98)
(21) International Application Number: PCT/EP97/05619		(81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, HU, ID, IL, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).
(22) International Filing Date: 11 October 1997 (11.10.97)		
(30) Priority Data: 96116373.0 12 October 1996 (12.10.96) EP		
(34) Countries for which the regional or international application was filed: DE et al.		
(71) Applicant (for all designated States except US): EVOTEC BIOSYSTEMS GMBH [DE/DE]; Grandweg 64, D-22529 Hamburg (DE).		
(72) Inventor; and (75) Inventor/Applicant (for US only): KASK, Peet [EE/EE]; Uus 5-3, EE3051 Harku (EE).		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.
(74) Agents: MEYERS, Hans-Wilhelm et al.; Postfach 10 22 41, D-50462 Köln (DE).		
(54) Title: METHOD OF ANALYSIS OF SAMPLES BY DETERMINATION OF A FUNCTION OF SPECIFIC BRIGHTNESS		
(57) Abstract		
<p>A method for characterizing samples having units, by monitoring fluctuating intensities of radiation emitted, scattered and/or reflected by said units in at least one measurement volume, the monitoring being performed by at least one detection means, said method comprising the steps of: a) measuring in a repetitive mode a number of photon counts per time interval of defined length, b) determining a function of the number of photon counts per said time interval, c) determining a function of specific brightness of said units on basis of said function of the number of photon counts.</p>		
<p>Mean count rate: ● 7.7 kHz ○ 193 kHz</p> <p>Number of events</p> <p>Number of counts</p> <p>A</p>		
<p>Weighted residuals</p> <p>Number of counts</p> <p>B</p>		

PATENT COOPERATION TREATY

PCT

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

Applicant's or agent's file reference ME/kk 972076wo	FOR FURTHER ACTION See Notification of Transmittal of International Preliminary Examination Report (PCT/IPEA/416)	
International application No. PCT/EP97/05619	International filing date (day/month/year) 11/10/1997	Priority date (day/month/year) 12/10/1996
International Patent Classification (IPC) or national classification and IPC G01N15/02		
Applicant AG EVOTEC BIOSYSTEMS GMBH et al.		


- This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.
- This REPORT consists of a total of 5 sheets, including this cover sheet.
 - ☒ This report is also accompanied by ANNEXES, i.e., sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

(3)

These annexes consist of a total of 5 sheets.

- This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☐ Priority
- III ☐ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- IV ☐ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☒ Certain defects in the international application
- VIII ☐ Certain observations on the international application

Date of submission of the demand 09/05/1998	Date of completion of this report 28.12.98
Name and mailing address of the IPEA/  European Patent Office D-80298 Munich Tel. (+49-89) 2399-0, Tx: 523656 epmu d Fax: (+49-89) 2399-4465	Authorized officer Strohmayer, B Telephone No. (+49-89) 2399-2669

**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT**

International application No. PCT/EP97/05619

I. Basis of the report

1. This report has been drawn on the basis of (*substitute sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to the report since they do not contain amendments.*):

Description, pages:

1-27 as originally filed

Claims, No.:

1-28 as received on 03/11/1998 with letter of 02/11/1998

Drawings, sheets:

1/12-12/12 as originally filed

2. The amendments have resulted in the cancellation of:

- ☐ the description, pages:
☐ the claims, Nos.:
☐ the drawings, sheets:

3. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

4. Additional observations, if necessary:

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/EP97/05619

V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

1. Statement

Novelty (N)	Yes:	Claims	1-28
	No:	Claims	
Inventive step (IS)	Yes:	Claims	
	No:	Claims	1-28
Industrial applicability (IA)	Yes:	Claims	1-28
	No:	Claims	

2. Citations and explanations

see separate sheet

VII. Certain defects in the international application

The following defects in the form or contents of the international application have been noted:

see separate sheet

Section V

D1=Quian and Elson : "Distribution of molecular aggregation by Analysis of fluctuation moments"

D3=EP601714

1. The subject matter of claim 1 is not inventive within the meaning of Art.33(3) PCT for the following reasons:

D1 (p.5479, abstract and first paragraph; p.5479, right col., 3rd and 5th paragraph, p.5482, left col. and right col., chapter "RESULTS", p.5483, last paragraph; p.5479, right col., lines 11-13 from bottom; Fig.3 and first two lines of the caption of Figure 3, the last seven lines of the caption of Fig.3 and p.5482, right column, lines 15-18 "The distributions ... Fig.3a" and the last paragraph of chapter "RESULTS") discloses a method for characterizing samples having particles, by monitoring fluctuating intensities of radiation emitted, scattered and/or reflected by said particles in at least one measurement volume, the monitoring being performed by at least one detection means, said method comprising the steps of

- a) measuring in a repetitive mode a number of photon counts per said time interval of defined length,
- b) determining a distribution function of the number of photon counts per said time interval,
- c) determine a distribution function of specific brightness of said particles on basis of said distribution function of the number of photon counts.

In D1 the distribution function of specific brightness indicates the mean number N_k in the effective illuminated volume v of particles emitting in the mean q_k photons per dwell time, where k denotes the various kinds of particles. The effective illuminated volume takes into account parameters of the spatial brightness function characteristic of the optical set-up (D1, p.5480, left col., l.20ff; compare appl., p.6, 1st paragraph).

The distribution function of specific brightness, i.e. the values N_k are calculated from the moments of the distribution function of the number of photon counts.

The subject matter of claim 1 differs from this prior art in that the distribution function of specific brightness of said particles is determined from the distribution function of the number of photon counts by finding a close fit between the experimentally determined

**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/EP97/05619

and a theoretical distribution function of the number of photon counts.

The obvious problem to be solved is to determine the distribution function of specific brightness of said particles directly from the distribution function of the photon counts by some sort of inversion. This problem is not new (l.5,6 of abstract; p.5479, left col.,l.21-24; p.5482, last paragraph of chapter "RESULTS").

In order to solve the problem, the skilled person is aware that the final result, i.e. the distribution function of the specific brightness is characterised by a limited number of parameters, which can be parameters of model distributions or discrete values of the distribution function itself. It is obvious for the skilled person that these parameters must be chosen to obtain the closest possible fit with the experimental data. It is to be noted that in the current application the term "fit" is used in a very broad meaning and includes any sort of inverse transformation (see for example p.12, last paragraph or dependent claim 21).

2. The subject matters of dependent claims 2-28 are likewise not inventive for the following reasons:

2: D1, p.5480, left col., l.24 ("optical sectioning")

3: D1, p.5480, left col., l.7-11 from bottom

4-6: D1, abstract

7-28: D1, see passages cited above with respect to claim 1 and additionally the chapter "Materials and Methods"; see also D3, Fig.2 and 12-18 and p.3,l.19-27, p.5,l.5-26, p.9,l.55-p.10,l.51 and p.12,l.3-12.

Section VII

The description is not in conformity with the claims. It contains for example several references to "units, preferably particles" (for example on p.4) or to the "function, preferably the distribution" (for example on p.5), although claim 1 is restricted to particles and the distribution function.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

Method of analysis of samples by determination
of a function of specific brightness

The present invention relates to a method for characterizing samples having units which emit, scatter and/or reflect radiation by measuring in a repetitive mode a number of photon counts per time interval of defined length and determining a function of the number of photon counts.

The first successful studies on fluorescence intensity fluctuations were performed by Magde, Elson and Webb (Biopolymers, Vol. 13, 29-61, 1974) who demonstrated the possibility to detect number fluctuations of fluorescent molecules and established a research field called fluorescence correlation spectroscopy (FCS). FCS was primarily developed as a method for determining chemical kinetic constants and diffusion coefficients. The experiment consists essentially in measuring the variation with time of the number of molecules of specific reactants in a defined open volume of solution. The concentration of a reactant is measured by its fluorescence from a small measurement volume. The measurement volume is defined by a focussed laser beam, which excites the fluorescence, and a pinhole in the image plane of the microscope collecting fluorescence. Intensity of fluorescence emission fluctuates in proportion with the changes in the number of fluorescent molecules as they diffuse into and out of the measurement volume and as they are created or eliminated by the chemical reactions. Technically, the direct outcome of an FCS experiment is the calculated autocorrelation function of the measured fluorescence intensity.

- 2 -

An important application of FCS is determination of concentrations of fluorescent species having different diffusion rates, in their mixture. In order to separate the two terms in the autocorrelation function of fluorescence intensity corresponding to translation diffusion of two kinds of particles, at least about a two-fold difference in diffusion time is needed, which corresponds to an eight-fold mass difference of particles. Furthermore, even if one succeeds in separating the two terms in the autocorrelation function of fluorescence intensity, it is yet not sufficient for determining the corresponding concentrations except if one knows the relative brightness of the two different types of particles.

Whereas conventional FCS yields rather limited information about aggregate sizes from a simple autocorrelation function of fluorescence intensity fluctuations, possible biophysical applications demand the ability to analyse complex mixtures of different species. For that purpose, Palmer and Thompson studied higher order correlation functions of fluorescence intensity fluctuations and have outlined methods for determining the number densities and relative molecular brightness of fluorescence of different fluorescent species (Biophys. J., Vol. 52, 257-270, August 1987). Their technique may in principle prove useful in detecting and characterizing aggregates of fluorescent-labeled biological molecules such as cell surface receptors, but has a major disadvantage of being rather complex, so that data processing of an experiment including the calculation of high-order correlation functions lasts hours.

A considerably less complicated method than calculation of high order auto-correlation functions for characterizing mixtures of fluorescent species of different specific brightness is calculation of higher order moments of fluorescence intensity out of the experimentally determined distribution of the number of photon counts. This method was presented by Qian and Elson (Biophys. J., Vol. 57, 375-380, February 1990; Proc. Natl. Acad. Sci. USA, Vol. 87, 5479-5483, July 1990). In their demonstration

- 3 -

experiments signal acquisition times of about 7 minutes were used for a relatively favourable experimental system of two kinds of fluorescent particles which differed by 30-fold in their specific brightness, the mixture of monomers and 30-mers. The method of moments is relatively simple and fast in calculations, but it allows to determine only a limited number of unknown parameters characterizing the sample because usually only about three or four first moments of fluorescence intensity can be calculated from the experiment with precision sufficient for further analysis.

Because of this reason, the method of moments is hardly suitable for characterizing complex samples or selecting between competing models of the sample or checking whether the given model is appropriate.

One object of the invention is to obtain reliable information about a sample having units emitting, scattering and/or reflecting photons, which renders possible an analysis of the sample with respect to certain ingredients or with respect to certain states of the sample.

Another object of the present invention is to substantially extend the useful information obtainable from the experimentally determined function, preferably distribution of the number of photon counts.

The objects of the present invention are solved with the method having the features of claim 1.

The term "unit of a sample" refers, in general, to subparts of the sample which are capable of emitting, scattering and/or reflecting radiation. A sample might contain a number of identical units or different units which preferably can be grouped into species. The term "different species" refers also to different states, in particular different conformational states, of a unit such as a molecule. Fluorescently labelled or natural-

ly fluorescent molecules, molecular complexes, vesicles, cells, beads and other particles in water or other liquids are examples of fluorescent units in liquid samples, while examples of fluorescent units of a solid sample are impurity molecules, atoms or ions, or other fluorescence centers.

What is meant by the term "specific brightness" of units in the sense of the present invention is a physical characteristic expressing in what extent a unit of given species is able to emit, scatter and/or reflect radiation, preferably light. It is thought to characterize single units, preferably particles, and therefore the value of specific brightness is not depending on concentration of the units, neither on the presence of other units. Thus, a change of the total count rate of photons emitted, scattered and/or reflected from the measurement volume, if only due to a change in concentration of the total number of units, does not influence the measured value of specific brightness and the value of the ratio of numbers of units of different species determined by the present invention. Specific brightness is usually expressed in terms of the mean count rate per unit which is a weighted average of the count rate over coordinates of the unit in the measurement volume. In some cases, one might prefer to express specific brightness in count rates corresponding to a unit positioned in a place where the count rate has its top values. This could e.g. be the center of the focus of an incident beam.

The importance of the present invention for the analysis of samples may be illustrated by the following, non-limiting example: Assuming that a solution contains a quantity (a) of one type of particles (A) with a respective specific brightness (I_a) and a quantity (b) of another type of particles (B) with a respective specific brightness (I_b), the overall count rate of photons emitted by the solution depends on the expression $I_a \cdot a + I_b \cdot b$. Thus, by mere determination of the overall count rate, it is not possible to dissolve the value of a and/or b. Generally, in fluorimetric measurements, the overall count rate

- 5 -

of at least one type of particles is determined in an independent experiment. If the total number $a+b$ of particles does not change with respect to this measurement, the ratio a/b or its inverse can be determined by mere determination of the overall count rate of the mixture in a second measurement. However, the assumption, that the total number $a+b$ does not change between the two measurements, is often wrong. For example, adsorption effects of particles to surfaces may occur. Fluorimetric measurements cannot verify the total number of particles $a+b$ independently. The present invention overcomes these restrictions. From one measurement, the numbers of particles a and b can be determined without any prior information of their respective specific brightnesses.

It is to be understood that the following description is intended to be illustrative and not restrictive. Many embodiments will be apparent to those of skill in the art upon reviewing the following description. By way of example, the invention will be described primarily with reference to measuring numbers of photon counts from light emitted by fluorescently labelled particles in a sample. For example, in some embodiments it may be desirable to measure numbers of photon counts of other origin than fluorescence.

The present invention provides a method for calculating the expected function, preferably the distribution of the number of photon counts corresponding to given real equipment and samples of given composition. The ability to predict the distribution of the number of counts corresponding to samples of given composition allows, when studying samples of unknown composition, to find out the model of the sample yielding the closest fit between the calculated and the experimentally determined distribution of the number of photon counts. What is meant by the composition of the sample here is the specific brightness and concentration of units present in the sample. For example, a solution of a single fluorescent dye is characterized by two parameters: the concentration and specific brightness of the

dye molecules. A mixture of two fluorescent dyes is characterized by four parameters: the concentrations and specific brightnesses of the two kind of molecules. A complex mixture can be characterized by the distribution function of concentration versus specific brightness of molecules. Conveniently, concentration is expressed as the mean number of units per measurement volume, and specific brightness is expressed as the mean count rate per unit. Preferably said units are particles.

From the other side, the function, e.g. the distribution of the number of photon counts depends not only on composition of the sample, but also on equipment: first of all, on the spatial brightness function characteristic to the optical set-up, and on some characteristics of the detector like its dark count rate, its dead time and probability of afterpulsing. In the interest of a high quality of analysis, which is indicated by achieving a close fit between the experimentally determined and the calculated curves, it is preferred to characterize the equipment adequately.

Claim 1 does not cover the method described by Qian and Elson, as their method compares estimated and calculated moments of the distribution of light intensity, but not directly the distribution of the number of photon counts. Qian and Elson's teaching misses the teaching of the present invention.

According to the invention, a new quality of analysis of samples containing units, preferably particles, which emit, scatter and/or reflect radiation, preferably light, becomes possible. In a first step, a number of photon counts from radiation emitted, scattered and/or reflected by the units in the sample is measured per time interval of defined length in a repetitive mode. A series of different time intervals can also be used for a more complex analysis. In step 1, the number of photon counts is measured in a repetitive mode, i.e. the number of detected photons is counted preferably many times, repeating the procedure in a series of preferably consecutive time intervals, in

- 7 -

order to obtain statistically meaningful data. The length of the time interval is the duration of the time interval during which the number of photon counts is determined. The length of the time interval is usually expressed in microseconds or milliseconds or other units of time.

In a second step of the method according to the invention, a function, preferably a distribution of the number of photon counts per said time interval is determined, which means that it is determined how many times one has obtained a certain number of photon counts. The distribution is a function of the number of photon counts, expressing either the relative or absolute number of observed (or expected) events when a particular number of photon counts was (or is) obtained.

In a third step, the experimentally determined distribution of photon counts is analyzed directly without intermediate steps to obtain a function, preferably a distribution of specific brightnesses of the units in the sample.

It is preferred that at least one detection means monitors said number of photon counts. Any detector which is capable to detect radiation emitted, scattered and/or reflected by units of the sample may be used, said radiation arising preferably out of at least one measurement volume. Appropriate detection means such as an avalanche photo-diode, a photomultiplier or conventional photo-diodes are well known to those of skill in the art. It might also be preferred to use a multidetector consisting of a monolithic configuration of a plurality of detectors, especially if one wants to measure a set of samples in parallel as it is the case in miniaturized high throughput screening. It might further be preferred to use a two-dimensional multi-array detector.

In the sense of the present invention, particles are preferably luminescently labelled or unlabelled, preferably naturally luminescent, molecules or macromolecules, or dye molecules, molecu-

lar aggregates, complexes, vesicles, cells, viruses, bacteria, beads, or mixtures thereof.

The luminescence properties of the units can be varied by conjugating them with a specific luminophore via different linker molecules. It might be preferred to use polymeric linker molecules consisting of a varying number of equal or different monomers.

It may be advantageous to provide at least one of the units with a specific site to which an affinity substance having detectable properties will bind.

In a preferred embodiment, at least one of the units has a tag of histidine residues to which an affinity substance such as a chelate complex can bind. It might be preferred to use complexes of nickel-nitrilotriacetic acid (Ni-NTA) and a luminescence label as said affinity substance. In a further preferred embodiment, said complex contains two or more chelating groups and at least one luminescence label.

The luminescence properties of the units may also be varied by conjugating them with a first molecule, as e.g. biotin, which binds a luminescently labelled second molecule, as e.g. luminescently labelled avidin or streptavidin, or vice versa as it is described in detail in example 3.

Luminescence properties of a unit can also be changed by energy transfer. Energy absorbed by a donor is transferred upon close contact to a luminophore of an acceptor and subsequently emitted.

In a further preferred embodiment, the units, preferably particles, each carry a number of binding sites for luminescent particles. Luminescent particles can directly or via secondary molecules bind to these binding sites. Since highly luminescent particles are generated when many luminescent particles bind to the binding sites of the first particles, the method according to the invention is able to distinguish easily between particles with a large difference in luminescence intensity, so that even a small amount of bound luminescent particles can be measured in presence of an excess concentration of free

luminescent particles. This embodiment provides a new analysis of particles which do not carry a luminescent label by binding to a second particle which is luminescently labelled, but whose brightness does not change upon binding. A commercially very important application of this method is the measurement of fluorescently labelled antibodies binding to an antigen, while the antigen is binding to at least one of the multiple binding sites of the particle which is preferably a bead, or vice versa. The method can also be applied to other types of interactions such as nucleic acid hybridization or protein/nucleic acid interaction. The invention can also be applied for the analysis of distribution characteristics of said particles, such as for quality control and process control of polymers or oligomeric suspensions of particles. In addition, surface areas of particles can be analyzed as well as distributions of surface areas of particles.

In a preferred embodiment, one type of particle, subsequently denoted A, carries more than one binding site. Another, luminescent type of particles, subsequently denoted C, can bind (i) either directly to at least one of the binding sites of particle A, or (ii) binds to at least one binding site of a molecule B, which in turn binds to at least one of the binding sites of particle A. These bindings may be mediated either by naturally occurring binding sites on the particles, or mediated by introduction of specific binding sites to the particles A, B and/or C. Since in both cases more than one of the particles of type C may bind to particle A, the complex will emit more photons than free particles of type C. This embodiment provides a convenient way to measure binding of particles of type B to a particle of type C or A, although the particle of type B is not luminescently labelled.

In a further preferred embodiment, the measurement volume is only a part of the total volume of the sample and said units, preferably particles, are diffusing and/or being actively transported into and out of said measurement volume and/or the

- 10 -

sample is actively transported and/or optically scanned. If said units, e.g. fluorescent particles, are sufficiently small, then diffusion is fast enough for data acquisition from a great number of independent measurement volumes, and data acquisition using time averaging is nearly identical to ensemble averaging. However, if the characteristic time of diffusion is substantially longer than the time interval necessary for measuring fluorescence intensity (which is usually 10 to 50 μs), then active transport (flow or scanning) can considerably save time of data acquisition.

The measurement volumes can preferably be arranged on two-dimensional carriers, such as membranes or sheets having wells. Suitable carrier systems are e.g. described in WO 94/16313 (EVOTEC BioSystems GmbH). The measurement volumes might also be arranged in a linear way, as e.g. in a capillary system.

In fluorescence studies, it may be advantageous to take measures for reducing the background count rate, arising from Raman scattering in the solute material and dark count rate of the detector, with respect to the count rate per particle. In particular, it is in some cases preferred to use measurement volumes smaller than 100 μm^3 , more preferably about 1 μm^3 . Advantageously, the high signal to background count rate and the small optical measurement volume may be achieved by using at least one microscope objective, preferably with a numerical aperture ≥ 0.9 , in a confocal manner for both focussing an incident laser beam and collecting radiation, preferably light, emitted, scattered and/or reflected by units, preferably particles, in said samples. A confocal microscope set-up is preferably used which comprises at least one microscope objective, a dichroic mirror, a pin-hole in the image plane of said microscope objective, a detection means, a data acquisition means, and optionally means for scanning and/or actively transporting the sample. A suitable device is disclosed in WO 94/16313 (EVOTEC BioSystems GmbH). In a preferred embodiment the pin-hole might be replaced by an appropriate detector, as it is also

- 11 -

described in WO 94/16313. It might further be preferred to choose a working distance between the microscope objective and the measurement volume in such a way that background contributions are minimized. Preferably, the working distance should be smaller than 1000 μm .

In a preferred embodiment of the method, multiple photon excitation is used to excite a particle. Multiple photon excitation means that the sum, difference or any combination of wave frequencies of two, three or more photons is used for excitation of the secondary emission of the sample which can be e.g. luminescence or second order Raman scattering. Such an excitation scheme has an advantage in the sense that the excitation probability is not linearly dependent on excitation intensity, but on the second or higher power. Thus, the multiple photon excitation is mostly limited to the volume of the laser focus, whereas outside the laser focus no spurious excitation is generated. Appropriate laser sources of picosecond or subpicosecond pulses are well known to those of skill in the art. The present invention profits from such an excitation scheme in the sense that less background is generated compared to single photon excitation, and that there is no pinhole necessary to restrict the measurement volume. Thus, the pinhole diameter and its imaging on the detector do not enter as modelling parameters in the spatial brightness function any more.

In a further preferred embodiment, the measurement volume is restricted by the use of elements of near field microscopy. These can be used for focussing the excitation radiation of the units, and/or collecting the radiation emitted, scattered and/or reflected by the units. Near field optical microscopy means here that the light passes through an aperture with at least one of its dimensions being smaller than the wavelength of the light used and which is in direct contact to the measurement volume. The aperture may consist of an opaque layer with at least one hole of said diameter or at least one slit of appropriate width and/or a tapered glass fiber or wave guide with a tip diameter

- 12 -

of said width, optionally coated with an opaque layer outside. A suitable device is disclosed in WO 96/13744 and in the German patent 44 38 391 (EVOTEC BioSystems GmbH).

Another preferred embodiment combines near field optical microscopy for the excitation light path, and conventional optical microscopy for the emission light path, or vice versa. The present invention profits from such a realization in the sense that the size of the measurement volume is reduced compared to conventional confocal microscopy. Thus, the present invention can be used to measure higher particle concentrations as with other optical schemes.

A sample is usually characterized by values of concentration and specific brightness of one or more species of units, e.g. types of fluorescent particles. In cases when one or more of these values are known beforehand, the goal of analysis is to determine unknown values, either those of concentration, or specific brightness, or both.

Two alternative methods for selecting the model yielding a fit between the experimentally determined and calculated functions, preferably distributions of the number of photon counts can be used. In one embodiment, the well-known least squares fitting method, where the sample is described by a finite (usually small) number of parameters, can be employed. The purpose is to find values of the parameters yielding the closest fit between the experimental and the calculated curves. According to the invention, values of concentrations and/or specific brightnesses of a number of species of units, e.g. types of fluorescent particles, can be estimated. In a further embodiment, another general method called inverse transformation with linear regularization (ITR) can be employed. ITR describes the sample using a semi-continuous distribution function of units, preferably particles, versus their specific brightness, and searches for the closest fit demanding that the solution is a smooth function (For the method of ITR, see, e.g., W. H. Press,

S. A. Teukolsky, W. T. Vetterling, B. P. Flannery, Numerical recipes in C: the art of scientific computing, second edition, Cambridge University Press, 1992, p. 808). It might further be preferred to use an inverse transformation with constraints (ITC) or an inverse transformation with regularization and constraints (ITRC). Because of statistical errors and limited sizes of measured data, inverse transformation is often an ill-posed mathematical problem, characterized by wild oscillations in its outcome. ITR, ITC and ITRC stabilize the mathematical problem by looking for a "regular" (e.g. a smooth) or constrained solution, for example by minimizing the sum of squared deviations of statistical data and a function of the solution itself, penalizing "irregular", usually irreproducible structures in the outcome, or values having no physical meaning. An example for constraining is disallowing negative values for concentration.

In the following, the invention is further illustrated in a non-limiting manner. Particularly, it is described how the expected distribution of the number of photon counts is determined.

A preferably many times repeated step in the calculation of the probability distribution of the number of photon counts is calculation of the probability distribution of the number of photon counts emitted, scattered and/or reflected by single species from a spatial section of the measurement volume with a constant value of spatial brightness. It is well known that the probability distribution of the number of particles in an open volume is Poissonian. Also, if the number of particles inside the spatial section is given, the number of detected photons per sampling time interval is Poisson distributed. Consequently, the overall distribution of the number of photon counts emitted, scattered and/or reflected by single species from a spatial section of constant brightness and detected by an ideal detector is compound Poissonian.

- 14 -

As the next step, one may study the case in which the measurement volume is divided into a number of spatial sections of constant brightness. If the values of volumes and spatial brightnesses in each of the sections are known, the distribution of the number of photon counts corresponding to each section can be determined separately. All these distributions are compound Poissonian. Furthermore, if distributions of the number of photon counts for all sections were known, the overall distribution can be determined through convolutions, using the fact that the total number of counts is the sum of the number of counts originating from different sections of the measurement volume.

As the following step, one may study a mixture of species, e.g. mixtures of fluorescent particles having different values of specific brightness. Each spatial section of the measurement volume can be divided into a number of abstract subsections each containing only particles of a single species. A similar procedure can be applied now as described above for spatial sections of the measurement volume in order to determine the overall distribution of the number of photon counts.

An experimentally determined distribution of the number of photon counts is ruled not only by properties of the light beam, but is influenced also by nonideal properties of the photon detector. Stochastically, the dark counts of the detector behave in the same manner as photon counts from background light of constant intensity. Their contribution are photon counts of Poisson distribution. Also, the way how the dead time of the detector and its afterpulsing distort the distribution of photon counts are known from literature on photon statistics (see e.g. B. Saleh, Photoelectron Statistics, Springer, Berlin, 1978).

In summary, the expected distribution of the number of photon counts is determined, from one side, by characteristics of the sample (concentrations and specific brightnesses of fluorescent particles of different kind), and, from the other side, by

- 15 -

characteristics of the equipment (the sampling time interval, the spatial brightness function, the background count rate, the dead time and the afterpulsing probability of the detector).

In one embodiment, both the dead time and the afterpulsing probability of the detector are determined from experiments in which the distribution of the number of photon counts corresponding to light of constant intensity is determined. Correction for the dead time of the detector may be performed on the basis of a formula derived by Cantor and Teich (J. Opt. Soc. Am. 65, 786, 1975; see also B. Saleh, p. 272-277). Mathematics of afterpulsing is simple: each photon pulse can be followed by another (artificial) pulse; this happens usually with a constant probability.

According to the invention, it is preferred that the spatial brightness function is characterized using experiments on a single species of particles. For example, if the laser wavelength 514.5 nm is used, then a solution of Rhodamine 6G is a convenient sample which can be used for characterizing the spatial brightness function.

What characteristics of the spatial brightness function can be employed when determining the expected distribution of the number of counts are values of volumes of the sections of the measurement volume corresponding to a selected set of values of the spatial brightness. Typically, a set of twenty or thirty values of the spatial brightness positioned at a constant distance from each other in the logarithmic scale have been selected, covering two or three orders of magnitude. Contribution from the lower brightness areas can be accounted for by a single parameter, their relative contribution to fluorescence intensity. Intensity fluctuations of this light can be neglected. Because of the large number of the sections of the measurement volume, it would be less preferred to consider volumes corresponding to each of the sections as independent variables. It is convenient to consider them as variables depending on a

- 16 -

few other parameters, and determine the values of these parameters which yield the closest fit between the experimentally determined and the calculated distribution of the number of photon counts. Conveniently, a relatively simple model of the optical set-up is applied, which is not accounting for aberrations of the optics used, and which determines volumes of the sections of the measurement volume. For instance, the volumes of the sections depend on values of the convergence angle of the laser beam and the diameter of the pinhole. It might therefore be preferred to use the pinhole dimensions and the convergence angle of the incident laser beam as modelling parameters of the spatial brightness function.

Alternatively, simple mathematical expressions with formal parameters can be used instead of physical models for determining the volumes of spatial sections. The values of the formal parameters should preferably be selected in such a way that the closest fit between experimental and calculated distributions of the number of photon counts is achieved. Formal flexible expressions are advantageous because they yield a good fit between experimental and theoretical distributions of the number of photon counts. Secondly, calculations based on simple mathematical expressions are very fast compared to those based on physical models.

According to the invention, it may in some cases be preferred to select the length of the sampling time interval in such a way that in average more than one, preferably one to ten, counts per unit are yielded.

It may further be preferred that the length of the time interval is in average smaller than the characteristic correlation time of radiation intensity fluctuations.

If more than two unknown parameters of the sample have to be estimated, it is preferred to have no more than a few, preferably about one unit, preferably particle, per measurement volume.

- 17 -

It is preferred to have less than 10 units per measurement volumes to obtain good signal-to-noise ratios.

In one embodiment, at least one individual unit is statistically analyzed in terms of its specific brightness which may fluctuate or change non-stochastically.

The method of the present invention is particularly advantageous because information losses and distortions are kept minimal. Furthermore, a new quality attainable by the present invention is that the data processing depends less on a definite mathematical model of the sample compared to the other techniques which are state of the art. This means that the method is more reliable in terms of long term stability of an instrumental realization, and that any disturbance of the measurement volume by e.g. turbid samples or particles inside the laser beam does not significantly influence experimental results.

A further new quality attainable by the present invention is that the signal-to-noise ratio is much better compared to the techniques of the prior art. This means that experiments can be made within significantly shorter time (up to 100 fold shorter) than previously, showing the same signal-to-noise ratio as previous long term experiments. Since photo-bleaching of fluorescent molecules or fluorescently labelled molecules is an unsolved problem so far with any applied measurement technique in this field, especially when applied to cells, one is restricted to short measurement times. Thus, compared to the prior art, the present invention is advantageous for measurements inside cellular systems.

In one preferred embodiment, the present invention is realized in the field of fluorescence intensity fluctuation studies. The optical equipment is a conventional confocal FCS microscope equipped with a cw laser of visible light. The excitation laser beam is focussed into a sample which is a homogeneous water solution of a low concentration, typically in the nanomolar

- 18 -

range, of fluorescent material. Fluorescence emission from a microscopic confocal volume of about $1 \mu\text{m}^3$ is collected on a photon detector. The measurement time which is typically 1 to 60 seconds is divided into hundreds of thousands time intervals of typical width of 10 to 50 μs . The highest number of photon counts typically obtained in this experimental realization of the invention herein described is between 10 and 100.

The method according to the present invention is particularly well suited for use in high throughput screening, diagnostics, monitoring of polymerization, aggregation and degradation processes, or analytics of nucleic acids.

In screening procedures, substances that are possibly pharmacologically active can be analyzed through their interaction with specific receptors by examining said interaction with binding of a luminescently labelled ligand to receptors wherein natural receptors on their carrier cells as well as receptors on receptor-overexpressing carrier cells or receptors on vesicles or receptors in the form of expressed molecules or molecular complexes may be used. Moreover, the interaction of substances with enzymes in solution or in their genuine cellular environment can be detected by monitoring a change of the substrate of the enzyme, e.g. a change in brightness. Further applications, especially concerning the performance of assays, are disclosed in WO 94/16313 (EVOTEC BioSystems GmbH).

For the detection of specific recognition reactions, potential active substances can be present in complex natural, synthetic or semisynthetic mixtures which are subjected to separation prior to analysis. These mixtures can be separated first e.g. by chromatography to test the individual fractions for the presence of functional compounds preferably "on line" in a capillary at the end of a separation matrix. The coupling of fractionating methods with FCS detection is described in detail in WO 94/16313 (EVOTEC BioSystems GmbH). A similar set-up can be used with respect to the present invention.

- 19 -

Often, aggregation and degradation are phenomena to be monitored. Aggregates display brightnesses different from the monomers and can be monitored according to the present invention.

In sequencing according to the method of Sanger, oligomers of different length, of which the terminating nucleic acid is labelled with a dye, are identified. Advanced techniques, as e.g. the one described in DE 38 07 975 A1, use dyes which exhibit different properties, such as fluorescence lifetime, according to the type of base they are attached to. The determination of a base is much more secure if several properties, such as fluorescence lifetime and brightness, or any other specific physical property, are determined and cross checked for consistency. In a preferred embodiment, the sample to be sequenced is separated by gel or capillary electrophoresis, or a separation step is conducted by capillary electro-chromatography, electrohydrodynamic migration or related electrokinetic methods.

Example 1

The nature and advantages of the invention may be better understood on the basis of the following example where a mixture of rhodamine dyes is analyzed. Figures 1 to 7 illustrate consecutive steps of the analysis and their results.

Figure 1. Distributions of the number of photon counts experimentally determined at constant light intensities, time interval of 10 μ s and data collection time of 50 s. From curve fitting, the dead time of the detector was estimated to be 28 ± 4 ns; afterpulsing probability 0.0032 ± 0.0008 . In the lower graph, weighted residuals of the curve fitting are presented.

Figure 2. Distribution of the number of photon counts experimentally determined for a solution of rhodamine 6G at time interval of 40 μ s and data collection time of 50 s.

- 20 -

Figure 3. Normalized sizes of volumes of the twenty spatial sections of the measurement volume of the highest brightness. Section 0 corresponds to the maximal value of the spatial brightness while section 19 corresponds to about 100 times lower brightness. Sizes of volumes are determined from experiments on single fluorescent species.

Figure 4. Residuals of curve fitting corresponding to an experiment on rhodamine 6G solution (the experiment graphed by Figure 2).

Figure 5. Distribution of the number of photon counts experimentally determined for three samples at time interval of 40 μ s and data collection time of 50 s. The distribution corresponding to rhodamine 6G is the same as in Figure 2.

Figure 6. The results of the inverse transformation with linear regularization applied to the data of Figure 5.

Figure 7. Residuals corresponding to analysis of an experiment on the mixture solution of rhodamine 6G and tetramethylrhodamine (measured data in Figure 5). Graph a: expected curve was obtained by inverse transformation with linear regularization. Graph b: expected curve was obtained by the least squares fitting of the experimental data. Graph c: residuals of the least squares curve fitting under a wrong assumption of single species.

As the first preparatory step of analysis, the dead time and the afterpulsing probability of the photon detector are estimated. This was done by determining the distribution of the number of photon counts under illumination of the detector by light of constant intensity. Since the dead time distortions are most noticeable at high count rates while the afterpulsing distortions are better resolved at low count rates, the values of the dead time and the probability of afterpulsing were determined by jointly fitting distributions of the number of photon counts

- 21 -

determined at a relatively high and at a relatively low illumination intensity. The experimentally determined count number distributions are presented in Figure 1, together with residuals of the curve fitting. Values of the estimated parameters for the photon detector which have been used are: dead time 28 ns, afterpulsing probability 0.003.

The background count rate of the equipment is determined by measuring the count rate when the sample holder is filled with pure water.

As the second preparatory step, the spatial brightness distribution corresponding to a given optical set-up was characterized. For that purpose, the distribution of the number of photon counts corresponding to a solution of rhodamine 6G was experimentally determined (Figure 2). If the spatial brightness distribution is appropriately characterized, then the calculated curve fits the experimental curve. In order to numerically calculate the expected distribution of the number of photon counts, values of twenty one parameters characterizing the spatial profile are needed in our program: twenty sizes of volumes corresponding to twenty spatial sections of the highest values of spatial brightness, and the relative contribution to fluorescence light originating from areas of lower spatial brightness. In order to calculate values of these unknown parameters, a simple model of the optical equipment not accounting for aberrations was taken into use. As illustrated by Figure 3, the determined sizes of the twenty volumes are reproducible, even if other species than rhodamine 6G are used.

Having determined values of the twenty one parameters characterizing the spatial brightness distribution in the way just described above, the calculated distribution of the number of photon counts indeed fits the experimental curve, see Figure 4.

After the preparatory steps described above the equipment is ready for analysis. In Figure 5, distributions of the number

- 22 -

of photon counts corresponding to three different samples are presented. In Figure 6, the results of the inverse transformation with linear regularization are graphed. Both spectra corresponding to single species (rhodamine 6G or tetramethylrhodamine) have a single peak, but the peaks are centered at different values of specific brightness. The peak of rhodamine 6G is situated at about 108 kHz/molecule, whereas the peak of tetramethylrhodamine is centered at about 37 kHz/molecule. This indicates that a rhodamine 6G molecule is about 3 times brighter than a tetramethylrhodamine molecule. The spectrum corresponding to the mixture of the two species has two peaks centered indeed near the values obtained for the two species separately.

Figure 7 illustrates the residuals corresponding to the measurements of the mixture of rhodamine 6G and tetramethylrhodamine. Different methods of data processing yield slightly different fit curves (and different residuals). The upper graph corresponds to the spectrum of specific brightness obtained by inverse transformation with linear regularization. The middle graph corresponds to the fit curve obtained assuming two species. These two graphs are nearly identical. The experimentally determined distribution of the number of photon counts can formally be fitted under the wrong assumption of single species, which is shown in the lower graph, but the fit curve is obviously apart from the experimental one.

Example 2

To further demonstrate the usefulness of the present invention, a hybridization process was studied using a conventional confocal FCS microscope. A model system based on the interaction of two 32-base oligonucleotides, both labelled with the fluorescent dye TAMRA (5- (and 6-) carboxytetramethylrhodamine), has been investigated. The sequence of these oligonucleotides included a site for the restriction enzyme EcoRI enabling the cleavage of the primer dimer. This restriction analysis was used as a control in order to check the specificity of the results obtained by the method according to the present invention.

- 23 -

The hybridization was performed in a 10 mM Tris buffer (pH 7.4) containing 1 mM EDTA and 100 mM NaCl, the restriction analysis in 50 mM Tris-HCl (pH 7.5), 10 mM MgCl₂, 100 mM NaCl, 0.2% Triton X-100. The measurement time for each analysis was 30 seconds.

Figure 8. Number of particles per volume element as a function of their specific brightnesses.

The analysis of the TAMRA-labelled single-strand oligonucleotides revealed a single characteristic fluorescence intensity peak of 45 kHz for oligonucleotide A (Figure 8a), and of 20 kHz for oligonucleotide B (Figure 8b) upon excitation at 543 nm.

The hybridization of the oligonucleotides A and B resulted in a single intensity peak of 35 kHz (Figure 8c) indicating that the hybridization was complete. From the intensity values of the individual oligonucleotides, one would have expected an intensity peak of 65 kHz for this primer dimer. This discrepancy in the intensity can be explained by the occurrence of dye-nucleotide interactions and electron transfer-induced quenching.

Further evidence that the 35 kHz intensity peak represents the primer dimer is given by an cleavage experiment using EcoRI (Figure 8d). Restriction cleavage of the annealed primers results in an intensity peak of 20 kHz. The broadening of the intensity distribution indicates that the reaction was not complete.

These experiments demonstrate that the method according to the present invention is well-suited for studying hybridization processes which play an important role for the detection and characterization of pathogens. The ability of the method according to the present invention to measure the activity of a restriction endonuclease was also demonstrated by this series of experiments.

Example 3

Biotin labelled with the following different dyes

- a) 5- (and 6-) carboxy-X-rhodamine (abbr. ROX)
- b) 5- (and 6-) carboxytetramethylrhodamine (abbr. TAMRA)
- c) Rhodol Green™
- d) Rhodamin Green™
- e) Resorufine
- f) Texas Red® and
- g) Rhodamine B

with and without a spacer molecule (abbr. Sp.) as well as their mixtures with streptavidin have been analyzed according to the method of the present invention in order to monitor quenching effects of differently labelled biotin upon streptavidin binding.

Figure 9 shows the specific brightnesses of the different molecules. The presence of streptavidin is indicated by "+", whereas its absence is indicated by "-".

Example 4

The following example proves that the method according to the present invention is valid for the measurement of ligands bound to receptor populations on membrane vesicles.

The use of crude biological material such as biomembranes derived from tissue or cells in assays to be analyzed by single particle fluorescence detection analysis brings along issues dealing with the nature and heterogeneity of this material. When membrane preparations are generated from receptor-overexpressing cells, the most probable situation will be that there are many receptor molecules on a single membrane vesicle. For an analysis of fluorescent ligand binding to those receptors, the method according to the present invention is ideally suited because it discriminates between particles which display different fluorescence intensities. Membrane vesicles are slowly moving

- 25 -

particles (mean diffusion time $\tau_{diff} > 10$ ms) in a low concentration. Thus to attain a reasonable signal-to-noise ratio, the measurement times for these rare events have to be prolonged compared to nanomolar fluorophore solutions. In order to shorten measurement time and improve statistical accuracy, the effective volume to be analyzed had to be increased substantially without losing the advantage of detecting single molecules. This has been achieved by introducing a beam scanning device. Using a beam scanner also circumvents bleaching effects since the mean excitation time of a single vesicle is minimized by the movement of the laser beam.

The feasibility to quantify a biological interaction by the method according to the present invention is demonstrated using epidermal growth factor (abbr. EGF) binding to membrane vesicles from A431 human carcinoma cells. These cells express 10^5 to 10^6 epidermal growth factor receptors per cell.

Membrane vesicle preparations

Membrane preparations were carried out by cell disruption in a hypotonic buffer (20 mM Tris/HCl, pH 7.5, 5 mM $MgCl_2$) containing protease inhibitors (leupeptin, aprotinin and PMSF) using a glass homogenizer and high-spin centrifugation after removal of nuclei at low g force. 10% sucrose was added during the first centrifugation step. The membranes were homogenized in EGF binding buffer (20 mM HEPES pH 7.4, 140 mM NaCl, 5 mM $MgCl_2$, 1.8 mM CaCl, 4.2 mM $NaHCO_3$ and 5.5 mM glucose) using a Branson sonifier prior to the experiment. Protein content was determined with bicinchoninic acid (PIERCE) to 0.504 mg/ml (A431).

EGF binding studies

Binding experiments using EGF labelled with tetramethylrhodamine (abbr. TMR) and A431 membranes were performed according to Carraway et al. (J. Biol. Chem. 264:8699, 1989). Briefly, they were diluted with EGF binding buffer and were incubated with labelled ligand in 20 μ l samples for 40 minutes at room temperature. In competition experiments, membranes were incubated with

unlabelled EGF. Measurements of 30 seconds duration were carried out using one-dimensional beam scanning at 25 Hz and an amplitude of 700 μm .

Figure 10 shows a plot of the amount of ligand-receptor complexes in relation to the total ligand concentration.

Figure 11 shows examples of intensity distributions measured at certain concentrations of A431 vesicles.

The y-axis of the intensity distributions in figure 11 is constructed by multiplying the particle number obtained for each intensity in the grid of intensities by the intensity of that grid point, thus representing the contribution of particles at that intensity to the total intensity. It is preferred to choose this transformation as it emphasizes on particles with high intensity, but low concentration which is the case for vesicles with bound ligand. In terms of particle numbers (concentration), vesicles would only make a negligible contribution.

Figure 11a shows the intensity distribution of the ligand alone. The ligand has an intensity of about 40 kHz/particle. In figures 11 b-d, intensity distributions with increasing concentrations of vesicles are plotted. The increase of fluorescence from bright particles, i.e. vesicles with many ligand-receptor complexes on them, is clearly visible.

In order to quantify the degree of binding, one has to distinguish between unbound ligand and ligand-receptor complexes. Vesicles with several bound ligands are brighter than the ligand alone. Thus, a certain discriminating intensity I_d is chosen. Particles detected below this intensity are assumed to be ligand molecules, while particles detected above this intensity are counted as vesicles with receptor-ligand complexes.

The concentration c_l of free ligand is determined by summing up over all concentrations below the discriminating intensity:

- 27 -

$$C_L = \sum_{\forall i: I_i < I_d} C_i$$

The concentration c_{RL} ligand-receptor complexes is given by the assumption that a vesicle with n bound ligands has an intensity of n times the ligand intensity. Thus, an intensity component at an intensity I is from a vesicle with $n = I / I_{Ligand}$ ligand-receptor complexes, and the concentration of these complexes is given by

$$C_{RL} = \sum_{\forall i: I_i \geq I_d} C_i \cdot \frac{I_i}{I_{Ligand}}$$

Now, the degree of complex formation is given by

$$complex = \frac{C_{RL}}{C_L + C_{RL}}$$

This is plotted in figure 10 for the binding of labelled EGF to A431 vesicles.

Claims

1. A method for characterizing samples having units, by monitoring fluctuating intensities of radiation emitted, scattered and/or reflected by said units in at least one measurement volume, the monitoring being performed by at least one detection means, said method comprising the steps of:
 - a) measuring in a repetitive mode a number of photon counts per time interval of defined length,
 - b) determining a function of the number of photon counts per said time interval,
 - c) determining a function of specific brightness of said units on basis of said function of the number of photon counts.
2. A method according to claim 1, wherein said function of the number of photon counts per said time interval and/or said function of specific brightness is a distribution function.
3. The method according to claim 1 and/or 2, wherein said units are molecules, macromolecules, dyes, molecular aggregates, complexes, vesicles, cells, viruses, bacteria, beads, centers, or mixtures thereof in solids, liquids or gases.
4. The method according to at least one of the claims 1 to 3, wherein said units can be grouped into species which can be distinguished by their specific brightness.
5. A method according to at least one of the claims 1 to 4, wherein at least one species is luminescent, preferably fluorescent, and/or is luminescently labelled.

- 29 -

6. A method according to at least one of the claims 1 to 5, wherein the luminescence properties of the units are varied by conjugating them with a first molecule, in particular biotin, which binds a luminescently labelled second molecule, in particular luminescently labelled avidin or streptavidin, or vice versa.
7. A method according to claim 6 wherein the first molecule is a (6xHis)tag and the second molecule is a luminescently labelled Ni-NTA-derivatives.
8. A method according to at least one of the claims 1 to 7, wherein the luminescence properties of a unit are changed by energy transfer, in which energy absorbed by said unit is transferred upon close contact to a luminophore of an acceptor and subsequently emitted.
9. A method according to at least one of the claims 1 to 8, wherein said units each carry a number of binding sites for luminescent units.
10. A method according to at least one of the claims 1 to 9, wherein the measurement volume is only a part of the total volume of the sample and has a volume $\leq 10^{-12}$ l, preferably $\leq 10^{-14}$ l.
11. A method according to at least one of the claims 1 to 10, wherein said units are diffusing and/or being actively transported into and out of said measurement volume and/or the sample is actively transported and/or optically scanned.
12. A method according to at least one of the claims 1 to 11, wherein the measurement volumes are arranged on a two-dimensional carrier, in particular on a membrane or in sheets having wells, or in linear way, preferably in a capillary system.

- 30 -

13. A method according to at least one of the claims 1 to 12, wherein a confocal microscope set-up is used, comprising at least one microscope objective, preferably with a numerical aperture ≥ 0.9 , for both focussing an incident laser beam and collecting radiation emitted, scattered and/or reflected by said units of said sample, a dichroic mirror, a pin-hole in the image plane of said microscope objective, a detection means, a data acquisition means, and optionally means for scanning and/or actively transporting said sample.
14. A method according to at least one of the claims 1 to 13, wherein said measurement volume is restricted by the use of elements of near field optical microscopy, or their combination with conventional microscopy optics.
15. A method according to at least one of the claims 1 to 14, wherein fluorescence is induced using multiple photon excitation.
16. A method according to at least one of the claims 1 to 15, wherein the parameters of a spatial brightness function characteristic for the optical set-up are determined by measuring numbers of photon counts per defined time intervals in a repetitive mode from radiation emitted, scattered and/ or reflected by a single species.
17. A method according to at least one of the claims 1 to 16, wherein the dimension and two-dimensional shape of the pinhole positioned in the focal plane of the microscope is used as a modelling parameter of the spatial brightness function.
18. A method according to at least one of the claims 1 to 17, wherein the convergence angle of the incident laser beam is used as a modelling parameter of the spatial brightness function.

- 31 -

19. A method according to at least one of the claims 1 to 18, wherein the concentration and/or specific brightness of at least one species of said units is determined.
20. A method according to at least one of the claims 1 to 19, wherein said distribution of the number of photon counts is fitted using a priori information on the sample.
21. A method according to at least one of the claims 1 to 20, wherein said distribution of the number of photon counts is processed by applying an inverse transformation with linear regularization and/or constraints.
22. A method according to at least one of the claims 1 to 21, wherein the experimental parameters of the detection means, in particular dead time and afterpulsing probability of the detection means, are determined by measuring number of photon counts per defined time interval in a repetitive mode while the detection means is exposed to light of constant intensity or high frequency laser pulses.
23. A method according to at least one of the claims 1 to 22, wherein background count rate of the equipment is determined.
24. A method according to at least one of the claims 1 to 23, wherein the length of said time interval is in average smaller than the characteristic correlation time of radiation intensity fluctuations.
25. A method according to at least one of the claims 1 to 24, wherein the length of said time interval is selected to yield in average more than one, preferably one to ten, photon counts per said unit.
26. A method according to at least one of the claims 1 to 25, wherein the concentration of the sample or the size of the

- 32 -

measuring volume is selected to have in average no more than a few, preferably about one unit per measurement volume.

27. A method according to at least one of the claims 1 to 26, wherein at least one individual unit is statistically analyzed in terms of its specific brightness which may fluctuate or change non-stochastically.
28. A method according to at least one of the claims 1 to 27 for use in high throughput screening, diagnostics, monitoring of polymerization, aggregation and degradation processes, or analytics of nucleic acids.




Fig. 1a




Fig. 1b




Fig. 2




Fig. 3




Fig. 4







Fig. 5



7/12




Fig. 7a




Fig. 7b




Fig. 7c

Number of counts

8/12

oligo A




Fig. 8a

oligo B






Fig. 8b

9/12

oligo A&B hybrid



oligo A&B hybrid + EcoR1



10/12

Biotin-Streptavidin Binding




Fig. 9

11/12




Fig. 10

12/12

Fig. 11a




Fig. 11b




Fig. 11c





Fig. 11d



INTERNATIONAL SEARCH REPORT

National Application No

PCT/EP 97/05619

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 G01N15/02

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	H. QIAN ET AL.: "Distribution of molecular aggregation by analysis of fluctuation moments" PROC. NATL. ACAD. SCI., vol. 87, July 1990, USA, pages 5479-5483, XP000647943 cited in the application see page 5479, column 1, paragraph 1 see page 5479, column 2, paragraph 3 see figure 3 see page 5483, column 1, paragraph 2 --- -/--	1

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

16 February 1998

Date of mailing of the international search report

24/02/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Zinngrebe, U

INTERNATIONAL SEARCH REPORT

National Application No
PCT/EP 97/05619

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>SM. NIE ET AL.: "real-time detection of single-molecules in solution by confocal fluorescence microscopy" ANALYTICAL CHEMISTRY, vol. 67, no. 17, 1 September 1995, COLUMBUS US, pages 2849-2857, XP000647270 see figure 3</p> <p style="text-align: center;">---</p>	1
A	<p>EP 0 601 714 A (HAMAMATSU PHOTONICS KK) 15 June 1994 see page 3, line 19-27 see page 5, line 5-26 see page 9, line 55 - page 10, line 51 see page 12, line 3-21</p> <p style="text-align: center;">---</p>	1
A	<p>US 4 788 443 A (FURUYA YOSHIYUKI) 29 November 1988 see column 1, line 5-23 see column 3, line 30 - column 4 see figures 1,4</p> <p style="text-align: center;">---</p>	1
A	<p>US 4 768 879 A (MCLACHLAN RICHARD D ET AL) 6 September 1988 see abstract see column 4, line 59 - column 5, line 15; figure 2</p> <p style="text-align: center;">-----</p>	1

INTERNATIONAL SEARCH REPORT

Information on patent family members

I. International Application No

PCT/EP 97/05619

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0601714 A	15-06-94	JP 2575270 B JP 6148076 A US 5528046 A	22-01-97 27-05-94 18-06-96
US 4788443 A	29-11-88	JP 1878140 C JP 6003413 B JP 62287130 A	07-10-94 12-01-94 14-12-87
US 4768879 A	06-09-88	NONE	

CLAIMS

1. A method for characterizing samples having particles, by monitoring fluctuating intensities of radiation emitted, scattered and/or reflected by said particles in at least one measurement volume, the monitoring being performed by at least one detection means, said method comprising the steps of:
 - a) measuring in a repetitive mode a number of photon counts per time interval of defined length,
 - b) determining a distribution function of the number of photon counts per said time interval,
 - c) determining a distribution function of specific brightness of said particles on basis of said distribution function of the number of photon counts by finding a close fit between the experimentally determined and a theoretical distribution function of the number of photon counts, the latter of which takes into account parameters of the spatial brightness function characteristic of the optical set-up.
2. The method according to claim 1, wherein values of volumes of the sections of the measurement volume corresponding to a selected set of values of the spatial brightness serve as a means to characterize said spatial brightness function of the optical set-up.
3. The method according to claim 1 and/or 2, wherein the parameters of the spatial brightness function characteristic for the optical set-up are determined by

measuring numbers of photon counts per defined time intervals in a repetitive mode from radiation emitted, scattered and/or reflected by a single species.

4. The method according to at least one of the claims 1 to 3, wherein said particles are molecules, macromolecules, dyes, molecular aggregates, complexes, vesicles, cells, viruses, bacteria, beads, or mixtures thereof in liquids or gases.
5. The method according to at least one of the claims 1 to 4, wherein said particles can be grouped into species which can be distinguished by their specific brightness.
6. The method according to at least one of the claims 1 to 5, wherein at least one species is luminescent, preferably fluorescent, and/or is luminescently labelled.
7. The method according to at least one of the claims 1 to 6, wherein the luminescence properties of the particles are varied by conjugating them with a first molecule, in particular biotin, which binds a luminescently labelled second molecule, in particular luminescently labelled avidin or streptavidin, or vice versa.
8. The method according to claim 7, wherein the first molecule is a (6xHis) tag and the second molecule is a luminescently labelled Ni-NTA-derivative.
9. The method according to at least one of the claims 1 to 8, wherein the luminescence properties of a particle are changed by energy transfer, in which energy absorbed by said particle is transferred upon close contact to a luminophore of an acceptor and subsequently emitted.

10. The method according to at least one of the claims 1 to 9, wherein said particles each carry a number of binding sites for luminescent particles.
11. The method according to at least one of the claims 1 to 10, wherein the measurement volume is only a part of the total volume of the sample and has a volume $\leq 10^{-12}$ l, preferably $\leq 10^{-14}$ l.
12. The method according to at least one of the claims 1 to 11, wherein said particles are diffusing and/or being actively transported into and out of said measurement volume and/or the sample is actively transported and/or optically scanned.
13. The method according to at least one of the claims 1 to 12, wherein the measurement volumes are arranged on a two-dimensional carrier, in particular on a membrane or in sheets having wells, or in linear way, preferably in a capillary system.
14. The method according to at least one of the claims 1 to 13, wherein a confocal microscope set-up is used, comprising at least one microscope objective, preferably with a numerical aperture ≥ 0.9 , for both focussing an incident laser beam and collecting radiation emitted, scattered and/or reflected by said particles of said sample, a dichroic mirror, a pin-hole in the image plane of said microscope objective, a detection means, a data acquisition means, and optionally means for scanning and/or actively transporting said sample.
15. The method according to at least one of the claims 1 to 14, wherein said measurement volume is restricted by the use of elements of near field optical microscopy, or their combination with conventional microscopy optics.

16. The method according to at least one of the claims 1 to 15, wherein fluorescence is induced using multiple photon excitation.
17. The method according to at least one of the claims 1 to 16, wherein the dimension and two-dimensional shape of the pinhole positioned in the focal plane of the microscope is used as a modelling parameter of the spatial brightness function.
18. The method according to at least one of the claims 1 to 17, wherein the convergence angle of the incident laser beam is used as a modelling parameter of the spatial brightness function.
19. The method according to at least one of the claims 1 to 18, wherein the concentration and/or specific brightness of at least one species of said particles is determined.
20. The method according to at least one of the claims 1 to 19, wherein said distribution of the number of photon counts is fitted using a priori information on the sample.
21. The method according to at least one of the claims 1 to 20, wherein said distribution of the number of photon counts is processed by applying an inverse transformation with linear regularization and/or constraints.
22. The method according to at least one of the claims 1 to 21, wherein the experimental parameters of the detection means, in particular dead time and afterpulsing probability of the detection means, are determined by measuring number of photon counts per defined time interval in a repetitive mode while the

detection means is exposed to light of constant intensity or high frequency laser pulses.

23. The method according to at least one of the claims 1 to 22, wherein background count rate of the equipment is determined.

24. The method according to at least one of the claims 1 to 23, wherein the length of said time interval is in average smaller than the characteristic correlation time of radiation intensity fluctuations.

25. The method according to at least one of the claims 1 to 24, wherein the length of said time interval is selected to yield in average more than one, preferably one to ten, photon counts per said particle.

26. The method according to at least one of the claims 1 to 25, wherein the concentration of the sample or the size of the measurement volume is selected to have in average no more than a few, preferably about one particle per measurement volume.

27. The method according to at least one of the claims 1 to 26, wherein at least one individual particle is statistically analyzed in terms of its specific brightness which may fluctuate or change non-stochastically.

28. The method according to at least one of the claims 1 to 27 for use in high throughput screening, diagnostics, monitoring of polymerization, aggregation and degradation processes, or analytics of nucleic acids.